

A Quick Tour of the EIC Physics Case


Marco Stratmann



marco@bnl.gov



this talks can only present some highlights of the EIC program



in the absence of the EIC white paper,
I refer to the **500+ pages** INT report
arXiv:1108.1713

presented results are
mainly due to the efforts of
the EIC task force at BNL
and other supporters of an EIC





1

inclusive deep-inelastic scattering in ep and eA

what to measure



bread and butter probe at an electron-ion collider



need to measure only the **scattered electron** (its **energy** and **angle**)

- fully determines **two relevant kinematical variables**
a 3rd variable (**inelasticity y**) is related to x, Q^2
through the available c.m.s. energy



Q^2 : virtuality of exchanged photon

x : momentum fraction of probed parton

what to measure



bread and butter probe at an electron-ion collider



need to measure only the **scattered electron** (its **energy** and **angle**)

- fully determines **two relevant kinematical variables**
a 3rd variable (**inelasticity y**) is related to x, Q^2
through the available c.m.s. energy

Q^2 : virtuality of exchanged photon
 x : momentum fraction of probed parton



beware: ☐ want to measure up to large Q^2 : exchange of W, Z bosons contributes

- need to determine x, Q^2 from hadronic final-state (lepton can turn into neutrino)

what to measure



bread and butter probe at an electron-ion collider



need to measure only the **scattered electron** (its **energy** and **angle**)

- fully determines **two relevant kinematical variables**
a 3rd variable (**inelasticity y**) is related to x, Q^2
through the available c.m.s. energy



Q^2 : virtuality of exchanged photon
 x : momentum fraction of probed parton

- beware:**
- ☐ want to measure up to large Q^2 : exchange of W, Z bosons contributes
 - need to determine x, Q^2 from hadronic final-state (lepton can turn into neutrino)
 - ☐ kinematics obscured by additional photon radiation off the lepton
 - need Monte Carlo tools to control; cannot be separated from detector acceptance

what to measure



bread and butter probe at an electron-ion collider



need to measure only the **scattered electron** (its **energy** and **angle**)

- fully determines **two relevant kinematical variables**
a 3rd variable (**inelasticity y**) is related to x, Q^2
through the available c.m.s. energy


Q^2 : virtuality of exchanged photon
 x : momentum fraction of probed parton

- beware:**
- ☐ want to measure up to large Q^2 : exchange of W, Z bosons contributes
 - need to determine x, Q^2 from hadronic final-state (lepton can turn into neutrino)
 - ☐ kinematics obscured by additional photon radiation off the lepton
 - need Monte Carlo tools to control; cannot be separated from detector acceptance
 - ☐ need to measure also the polarization (and luminosity) very well

why is this interesting and what can we learn?

DIS measurements are *the classic tool* to study the partonic structure of nucleons (nuclei):



- ☒ distribution of partons in longitudinal momentum x
- ☒ **gluon density** from studying Q^2 dependence for fixed x
- ☒ theory tools and higher order QCD corrections well understood
- ☐ not so good for separation of quark flavors
- ☐ no information on spatial distribution of partons



why is this interesting and what can we learn?

DIS measurements are *the classic tool* to study the partonic structure of nucleons (nuclei):



- ☒ distribution of partons in longitudinal momentum x
 - ☒ **gluon density** from studying Q^2 dependence for fixed x
 - ☒ theory tools and higher order QCD corrections well understood
 - ☐ not so good for separation of quark flavors
 - ☐ no information on spatial distribution of partons
- HERA has studied unpolarized ep scattering in great detail
 - discovered dramatic rise in gluon distribution
 - achieved 1-2 % uncertainties



why is this interesting and what can we learn?


DIS measurements are *the classic tool* to study the partonic structure of nucleons (nuclei):

- ☒ distribution of partons in longitudinal momentum x
 - ☒ **gluon density** from studying Q^2 dependence for fixed x
 - ☒ theory tools and higher order QCD corrections well understood
 - ☐ not so good for separation of quark flavors
 - ☐ no information on spatial distribution of partons
- HERA has studied unpolarized ep scattering in great detail
 - discovered dramatic rise in gluon distribution
 - achieved 1-2 % uncertainties
 - current data for **polarized ep and eA DIS** cover only a very restricted x, Q^2 range (no collider experiments so far)




extension of x, Q^2 coverage with an EIC

polarized ep scattering




eA scattering




extension of x, Q^2 coverage with an EIC

polarized ep scattering




eA scattering




- gain two decades in x -> get into the region where gluons and sea quarks dominate

extension of x, Q^2 coverage with an EIC

polarized ep scattering




eA scattering




- gain two decades in x -> get into the region where gluons and sea quarks dominate
- cover large Q^2 range for each x -> study “scaling violations” -> gluon density

extension of x, Q^2 coverage with an EIC


polarized ep scattering



eA scattering



- gain two decades in x -> get into the region where gluons and sea quarks dominate
- cover large Q^2 range for each x -> study "scaling violations" -> gluon density
- can reach large Q^2 (at medium-to-large x) -> access to electroweak effects




what can be achieved: spin structure

- RHIC will determine $\Delta g(x, Q^2)$ down to $x \approx \text{few} \times 10^{-2}$
but need access down to $\text{few} \times 10^{-4}$ to close chapter on spin
- an EIC can do just that ...

recall:



$$\Delta f(x) \equiv f_{\rightarrow\rightarrow}(x) - f_{\leftarrow\rightarrow}(x)$$



based on global QCD analyses with and without realistic EIC pseudo data


what can be achieved: spin structure

- RHIC will determine $\Delta g(x, Q^2)$ down to $x \approx \text{few} \times 10^{-2}$
but need access down to $\text{few} \times 10^{-4}$ to close chapter on spin
- an EIC can do just that ...

recall:



$$\Delta f(x) \equiv f_{\rightarrow\rightarrow}(x) - f_{\leftarrow\rightarrow}(x)$$



based on global QCD analyses with and without realistic EIC pseudo data


what can be achieved: spin structure

- RHIC will determine $\Delta g(x, Q^2)$ down to $x \approx \text{few} \times 10^{-2}$
but need access down to $\text{few} \times 10^{-4}$ to close chapter on spin
- an EIC can do just that ...

recall:



$$\Delta f(x) \equiv f_{\rightarrow\rightarrow}(x) - f_{\leftarrow\rightarrow}(x)$$



based on global QCD analyses with and without realistic EIC pseudo data


what can be achieved: spin structure

- RHIC will determine $\Delta g(x, Q^2)$ down to $x \approx \text{few} \times 10^{-2}$
but need access down to $\text{few} \times 10^{-4}$ to close chapter on spin
- an EIC can do just that ...


recall:



$$\Delta f(x) \equiv f_{\rightarrow}^{\rightarrow}(x) - f_{\leftarrow}^{\rightarrow}(x)$$



based on global QCD analyses with and without realistic EIC pseudo data




crucial impact on our understanding of decomposition of proton spin

what can be achieved: nuclear PDFs

goals:


- map the momentum distribution of sea quarks and gluons down to small x
- gather unambiguous evidence for non-linear QCD dynamics (aka **saturation**)



what can be achieved: nuclear PDFs

goals:

- map the momentum distribution of sea quarks and gluons down to small x
- gather unambiguous evidence for non-linear QCD dynamics (aka **saturation**)



key advantage of eA DIS over ep DIS

amplification of non-linear effects by nuclear “oomph factor”


$$Q_s^2(x, A) \sim \left(\frac{A}{x} \right)^{1/3}$$

expect to see saturation in eA at 100-200 × higher values of x than in ep

what can be achieved: nuclear PDFs

goals:

- map the momentum distribution of sea quarks and gluons down to small x
- gather unambiguous evidence for non-linear QCD dynamics (aka **saturation**)



key advantage of eA DIS over ep DIS

amplification of non-linear effects by nuclear "oomph factor"

$$Q_s^2(x, A) \sim \left(\frac{A}{x} \right)^{1/3}$$

expect to see saturation in eA at 100-200 \times higher values of x than in ep

- presence of (large enough) **saturation scale** Q_s allows one to perform quant. calculations in well-defined framework ("CGC")
- expect "physics at high gluon density" to be universal; can verify this at an EIC!

what can be achieved: nuclear PDFs

particularly suited inclusive observable: DIS structure function $F_L(x, Q^2)$ **unmeasured so far**

- extraction of F_L needs

“Rosenbluth separation”

measurements at fixed x, Q^2 for different y (i.e., S)

what can be achieved: nuclear PDFs


particularly suited inclusive observable: DIS structure function $F_L(x, Q^2)$ **unmeasured so far**

**expect strong non-linear effects in F_L
due to its sensitivity to gluons**

- extraction of F_L needs
“Rosenbluth separation”

measurements at fixed x, Q^2 for different y (i.e., S)

Au ($A=197$)



measure of non-linear effects



what can be achieved: nuclear PDFs

particularly suited inclusive observable: DIS structure function $F_L(x, Q^2)$ **unmeasured so far**

**expect strong non-linear effects in F_L
due to its sensitivity to gluons**


- extraction of F_L needs
“Rosenbluth separation”
measurements at fixed x, Q^2 for different y (i.e., S)
- can be done:

Au ($A=197$)



**combined analysis with other observables can
reveal presence of non-linear effects**

2




semi-inclusive probes in ep and eA

what to measure and why

extension of double-differential DIS cross section to a

(in general) **six-fold diff. cross section** $\frac{d\sigma}{dx dQ^2 dz d\phi_S d\phi_h dp_T^h}$




z: energy fraction of observed hadron

p_T^h: its transverse momentum

φ: azimuthal angle of hadron

φ_S: azimuthal angle of spin vector

what to measure and why



extension of double-differential DIS cross section to a

(in general) **six-fold diff. cross section** $\frac{d\sigma}{dx dQ^2 dz d\phi_S d\phi_h dp_T^h}$

z: energy fraction of observed hadron

p_T^h: its transverse momentum


φ: azimuthal angle of hadron

φ_S: azimuthal angle of spin vector

□ **p_T and φ integrated cross section** is simplest but versatile SIDIS observable

can be used for flavor separation (needs π's and K's) and determinations of fragmentation functions

what to measure and why



extension of double-differential DIS cross section to a

(in general) **six-fold diff. cross section** $\frac{d\sigma}{dx dQ^2 dz d\phi_S d\phi_h dp_T^h}$

z: energy fraction of observed hadron


p_T^h: its transverse momentum

φ: azimuthal angle of hadron

φ_S: azimuthal angle of spin vector

- **p_T and φ integrated cross section** is simplest but versatile SIDIS observable
 - can be used for flavor separation (needs π's and K's) and determinations of fragmentation functions
- **azimuthal modulations** lead to transverse momentum dependent PDFs (TMDs)

what to measure and why



extension of double-differential DIS cross section to a

(in general) **six-fold diff. cross section** $\frac{d\sigma}{dx dQ^2 dz d\phi_S d\phi_h dp_T^h}$

z: energy fraction of observed hadron


p_T^h: its transverse momentum

φ: azimuthal angle of hadron

φ_S: azimuthal angle of spin vector

- ❑ **p_T and φ integrated cross section** is simplest but versatile SIDIS observable
can be used for flavor separation (needs π's and K's) and determinations of fragmentation functions
- ❑ **azimuthal modulations** lead to transverse momentum dependent PDFs (TMDs)
- ❑ **hadron-hadron correlations** in eA give clean access to saturation physics

what to measure and why



extension of double-differential DIS cross section to a

(in general) **six-fold diff. cross section** $\frac{d\sigma}{dx dQ^2 dz d\phi_S d\phi_h dp_T^h}$

z: energy fraction of observed hadron

p_T^h: its transverse momentum


φ: azimuthal angle of hadron

φ_S: azimuthal angle of spin vector


- **p_T and φ integrated cross section** is simplest but versatile SIDIS observable
can be used for flavor separation (needs π's and K's) and determinations of fragmentation functions
- **azimuthal modulations** lead to transverse momentum dependent PDFs (TMDs)
- **hadron-hadron correlations** in eA give clean access to saturation physics

all SIDIS measurements require good particle ID in broad kinematic regime

1st example: flavor separation for helicity PDFs




1st example: flavor separation for helicity PDFs





physics of TMDs

- theoretically interesting multi-scale problem: Q^2, p_T^h
- TMD framework applicable for $Q^2 \gg p_T^h$
- studied only in fixed target regime
nothing known yet about sea quarks and gluons



physics of TMDs



- theoretically interesting multi-scale problem: Q^2, p_T^h
- TMD framework applicable for $Q^2 \gg p_T^h$
- studied only in fixed target regime
nothing known yet about sea quarks and gluons
- **yields momentum distribution of partons in the transverse plane**



anisotropy due to transverse
proton polarization in y direction

physics of TMDs

- theoretically interesting multi-scale problem: Q^2, p_T^h
- TMD framework applicable for $Q^2 \gg p_T^h$
- studied only in fixed target regime
nothing known yet about sea quarks and gluons
- **yields momentum distribution of partons in the transverse plane**




anisotropy due to transverse
proton polarization in y direction


- origin of certain TMDs deeply linked with color gauge invariance of QCD

physics of TMDs


- theoretically interesting multi-scale problem: Q^2, p_T^h
- TMD framework applicable for $Q^2 \gg p_T^h$
- studied only in fixed target regime
nothing known yet about sea quarks and gluons
- **yields momentum distribution of partons in the transverse plane**



- origin of certain TMDs deeply linked with color gauge invariance of QCD
- unintegrated gluon density has **connection to CGC physics at small x**




anisotropy due to transverse proton polarization in y direction



specific example: Sivers function

- slew of different TMDs can be measured by selecting certain ϕ modulations


example: Sivers function



specific example: Sivers function

- slew of different TMDs can be measured by selecting certain ϕ modulations

example: Sivers function




$$f_{q/P^\uparrow}(x, \mathbf{k}_\perp, S) = f_1(x, \mathbf{k}_\perp^2) - \frac{\mathbf{S} \cdot (\hat{\mathbf{P}} \times \mathbf{k}_\perp)}{M} f_{1T}^\perp(x, \mathbf{k}_\perp^2)$$

specific example: Sivers function

- slew of different TMDs can be measured by selecting certain ϕ modulations

example: Sivers function



$$f_{q/P^\uparrow}(x, \mathbf{k}_\perp, S) = f_1(x, \mathbf{k}_\perp^2) - \frac{\mathbf{S} \cdot (\hat{\mathbf{P}} \times \mathbf{k}_\perp)}{M} f_{1T}^\perp(x, \mathbf{k}_\perp^2)$$

unintegrated PDF




important link to physics of
gluon saturation at small x

specific example: Sivers function

- slew of different TMDs can be measured by selecting certain ϕ modulations

example: Sivers function




$$f_{q/P^\uparrow}(x, \mathbf{k}_\perp, S) = f_1(x, \mathbf{k}_\perp^2) - \frac{\mathbf{S} \cdot (\hat{\mathbf{P}} \times \mathbf{k}_\perp)}{M} f_{1T}^\perp(x, \mathbf{k}_\perp^2)$$

unintegrated PDF



important link to physics of
gluon saturation at small x




Sivers function

- measures spin-orbit correlations
- link to **parton orbital motion** (only through models)
- reveals non-trivial aspects of QCD color gauge invariance

specific example: Sivers function

- slew of different TMDs can be measured by selecting certain ϕ modulations

example: Sivers function




$$f_{q/P^\uparrow}(x, \mathbf{k}_\perp, S) = f_1(x, \mathbf{k}_\perp^2) - \frac{\mathbf{S} \cdot (\hat{\mathbf{P}} \times \mathbf{k}_\perp)}{M} f_{1T}^\perp(x, \mathbf{k}_\perp^2)$$

unintegrated PDF



important link to physics of
gluon saturation at small x




Sivers function

- measures spin-orbit correlations
- link to **parton orbital motion** (only through models)
- reveals non-trivial aspects of QCD color gauge invariance


Sivers asymmetry has been observed only in the valence quark regime

prospects for Sivers related measurements

extracted u-valence density




extracted u-sea density




prospects for Sivers related measurements

extracted u-valence density




extracted u-sea density



so far unmeasured **gluon Sivers fct**
can be probed in D-meson correlations

observable: azimuthal asymmetry
correlating the total k_T of the D-meson pair
with transverse spin of the nucleon



angle between proton spin and k_T of D-meson pair

probing saturation with di-hadron correlations

- corresponding measurement in dAu at RHIC one of the best hints for saturation right now
- much cleaner probe in eA: no spectator background from electron side

tests universality of high gluon density framework between eA and pA

probing saturation with di-hadron correlations

- corresponding measurement in dAu at RHIC one of the best hints for saturation right now
- much cleaner probe in eA: no spectator background from electron side

tests universality of high gluon density framework between eA and pA

observable: **azimuthal correlation of hadron pair**


expectation: back-to-back peak washed out
by multiple gluon interactions in CGC;
peak persists w/o saturation for ep \rightarrow eA

probing saturation with di-hadron correlations

- corresponding measurement in dAu at RHIC one of the best hints for saturation right now
- much cleaner probe in eA: no spectator background from electron side

tests universality of high gluon density framework between eA and pA

CGC expectation



observable: **azimuthal correlation of hadron pair**

expectation: back-to-back peak washed out by multiple gluon interactions in CGC; peak persists w/o saturation for ep \rightarrow eA

kinematics:

$$p_T^{\text{trig}} > 2 \text{ GeV}$$


$$1 \text{ GeV} \leq p_T^{\text{assoc}} \leq p_T^{\text{trig}}$$

probing saturation with di-hadron correlations

- corresponding measurement in dAu at RHIC one of the best hints for saturation right now
- much cleaner probe in eA: no spectator background from electron side

tests universality of high gluon density framework between eA and pA

CGC expectation




kinematics:

$$p_T^{\text{trig}} > 2 \text{ GeV}$$


$$1 \text{ GeV} \leq p_T^{\text{assoc}} \leq p_T^{\text{trig}}$$

observable: **azimuthal correlation of hadron pair**

expectation: back-to-back peak washed out by multiple gluon interactions in CGC; peak persists w/o saturation for ep \rightarrow eA




3



diffractive / exclusive processes

what to measure

- one of the surprises at HERA: large fraction of diffractive events (15% of total DIS rate)



close relative of DIS

need in addition


t : momentum transfer squared
 M_X : mass of diffractive final-state

variables can be traded for β and x_P
where $x_{Bj} = \beta x_P$

diffractive event characterized by large **rapidity gap** (angular region w/o particle flow)
mediated by **color neutral exchange** (e.g. colorless combination of 2 or more gluons)

what to measure

- one of the surprises at HERA: large fraction of diffractive events (15% of total DIS rate)



close relative of DIS

need in addition

t : momentum transfer squared
 M_X : mass of diffractive final-state

variables can be traded for β and x_P
where $x_{Bj} = \beta x_P$

diffractive event characterized by large **rapidity gap** (angular region w/o particle flow)
mediated by **color neutral exchange** (e.g. colorless combination of 2 or more gluons)

terminology:

coherent


proton / heavy nucleus stays intact

incoherent

proton / heavy nucleus breaks up

what to measure

- one of the surprises at HERA: large fraction of diffractive events (15% of total DIS rate)



close relative of DIS

need in addition

t : momentum transfer squared
 M_X : mass of diffractive final-state

variables can be traded for β and x_P
where $x_{Bj} = \beta x_P$

diffractive event characterized by large **rapidity gap** (angular region w/o particle flow)
mediated by **color neutral exchange** (e.g. colorless combination of 2 or more gluons)

terminology:

coherent

incoherent

proton / heavy nucleus stays intact

proton / heavy nucleus breaks up

- ep: detect intact protons in forward detectors


critical: IR design

- eA: need to tag on emitted neutrons from nuclear breakup (shown to be possible with near 100% efficiency)

diffractive physics - why relevant?

recall: diffractive pattern in optics

position of minima θ_i related to size R of screen



diffractive physics - why relevant?



recall: diffractive pattern in optics

position of minima θ_i related to size R of screen

similarly: in coherent (elastic) scattering

$d\sigma/dt$ resembles diffractive pattern

where $|t| \approx k^2 \theta^2$



diffractive physics - why relevant?


recall: diffractive pattern in optics

position of minima θ_i related to size R of screen

similarly: in coherent (elastic) scattering


$d\sigma/dt$ resembles diffractive pattern

where $|t| \approx k^2 \theta^2$



crucial differences:

- target not always "black disc"
→ sensitivity to "size" of probe / onset of black disc limit
- incoherent (inelastic) contribution



diffractive physics - why relevant?


recall: diffractive pattern in optics

position of minima θ_i related to size R of screen

similarly: in coherent (elastic) scattering

$d\sigma/dt$ resembles diffractive pattern


where $|t| \approx k^2 \theta^2$



crucial differences:

- target not always "black disc"
→ sensitivity to "size" of probe / onset of black disc limit
- incoherent (inelastic) contribution


strong sensitivity to gluons



$$d\sigma \sim [g(x)]^2$$

due to required
color-neutral exchange

particularly sensitive to saturation



diffractive physics - why relevant?


recall: diffractive pattern in optics

position of minima θ_i related to size R of screen

similarly: in coherent (elastic) scattering

$d\sigma/dt$ resembles diffractive pattern


where $|t| \approx k^2 \theta^2$



crucial differences:

- target not always “black disc”
→ sensitivity to “size” of probe / onset of black disc limit
- incoherent (inelastic) contribution

strong sensitivity to gluons




$$d\sigma \sim [g(x)]^2$$

due to required
color-neutral exchange

particularly sensitive to saturation

bonus:



$\frac{d\sigma}{dt}$ \longleftrightarrow Fourier transform \longleftrightarrow spatial distribution of gluons



ratio of diffractive to total cross section

- black disc limit characterized by $\sigma_{\text{diff}}/\sigma_{\text{tot}} = 1/2$ (recall: HERA sees $\approx 1/7$ in ep)
 \rightarrow large fraction of diffractive event is unambiguous signature for reaching the saturated limit

estimates for fraction of low-mass coherent diffraction in ep and eA at EIC kinematics:





- find:
- w/o non-linear effects eA/ep ratio stays roughly one
 - non-linear effects enhance σ_{diff} in eA scattering

ratio of diffractive to total cross section

- black disc limit characterized by $\sigma_{\text{diff}}/\sigma_{\text{tot}} = 1/2$ (recall: HERA sees $\approx 1/7$ in ep)
 \rightarrow large fraction of diffractive event is unambiguous signature for reaching the saturated limit

estimates for fraction of low-mass coherent diffraction in ep and eA at EIC kinematics:



find:

- w/o non-linear effects eA/ep ratio stays roughly one
- non-linear effects enhance σ_{diff} in eA scattering


day-1 signature for
saturation at an EIC

exclusive vector meson production

- **unique probe** - allows to measure momentum transfer t in eA diffraction

$$t = (\mathbf{p}_A - \mathbf{p}_{A'})^2 = (\mathbf{p}_{VM} + \mathbf{p}_{e'} - \mathbf{p}_e)^2$$

in general, one cannot detect the outgoing nucleus and its momentum





exclusive vector meson production

- **unique probe** - allows to measure momentum transfer t in eA diffraction

$$t = (\mathbf{p}_A - \mathbf{p}_{A'})^2 = (\mathbf{p}_{VM} + \mathbf{p}_{e'} - \mathbf{p}_e)^2$$

in general, one cannot detect the outgoing nucleus and its momentum



cross section strongly depends on **overlap with VM wave function**



- **small size (J/ψ)**
cuts off saturation region in dipole amplitude
- **large size (ϕ, ρ, \dots)**
“sees more of dipole amplitude”
→ more sensitive to saturation

exclusive vector meson production

- **unique probe** - allows to measure momentum transfer t in eA diffraction

$$t = (\mathbf{p}_A - \mathbf{p}_{A'})^2 = (\mathbf{p}_{VM} + \mathbf{p}_{e'} - \mathbf{p}_e)^2$$

in general, one cannot detect the outgoing nucleus and its momentum



cross section strongly depends on **overlap with VM wave function**

- **small size (J/ψ)**
cuts off saturation region in dipole amplitude
- **large size (ϕ, ρ, \dots)**
“sees more of dipole amplitude”
→ more sensitive to saturation

Q^2 variation controls size of probe
→ go in (small Q^2) and out (large Q^2) of saturation region



spatial distribution of gluons through diffraction

goal: going after the source distribution of gluons through **Fourier transform of $d\sigma/dt$**

spatial distribution of gluons through diffraction

goal: going after the source distribution of gluons through **Fourier transform of $d\sigma/dt$**

- find:**
- typical diffractive pattern for coherent (non-breakup) part
 - as expected, J/ψ less sensitive to saturation effects than larger ϕ meson




how does the imaging work - what do we learn?

idea: momentum transfer t conjugate to transverse position (impact parameter b)

→ expect small t relevant for large b and vice versa

- coherent part probes “shape of black disc”
- incoherent part (dominant at large t) sensitive to “lumpiness” of the source (fluctuations, hot spots, ...)

→ impact on our understanding of initial conditions of heavy ion collisions




how does the imaging work - what do we learn?

idea: momentum transfer t conjugate to transverse position (impact parameter b)

→ expect small t relevant for large b and vice versa

- coherent part probes “shape of black disc”
- incoherent part (dominant at large t) sensitive to “lumpiness” of the source (fluctuations, hot spots, ...)

→ impact on our understanding of initial conditions of heavy ion collisions



what you need to do:
$$\mathbf{F}(\mathbf{b}) \sim \frac{1}{2\pi} \int_0^\infty d\Delta \, \Delta \, J_0(\Delta b) \sqrt{\frac{d\sigma}{dt}} \quad t \simeq -\Delta^2$$

“amplitude”


how does the imaging work - what do we learn?

idea: momentum transfer t conjugate to transverse position (impact parameter b)

→ expect small t relevant for large b and vice versa

- coherent part probes “shape of black disc”
- incoherent part (dominant at large t) sensitive to “lumpiness” of the source (fluctuations, hot spots, ...)

→ impact on our understanding of initial conditions of heavy ion collisions



what you need to do:
$$\mathbf{F}(\mathbf{b}) \sim \frac{1}{2\pi} \int_0^\infty d\Delta \Delta \mathbf{J}_0(\Delta \mathbf{b}) \sqrt{\frac{d\sigma}{dt}} \quad t \simeq -\Delta^2$$

“amplitude”

challenges:


- coherent part drops fast with t ; statistical errors become large beyond $\sim 4^{\text{th}}$ minima
- accessible t range rather small ($|t| \lesssim 0.2 \text{ GeV}^2$) for coherent part

how does the imaging work - what do we learn?

idea: momentum transfer t conjugate to transverse position (impact parameter b)

→ expect small t relevant for large b and vice versa

- coherent part probes “shape of black disc”
- incoherent part (dominant at large t) sensitive to “lumpiness” of the source (fluctuations, hot spots, ...)



→ impact on our understanding of initial conditions of heavy ion collisions

what you need to do:
$$\mathbf{F}(\mathbf{b}) \sim \frac{1}{2\pi} \int_0^\infty d\Delta \Delta \mathbf{J}_0(\Delta \mathbf{b}) \sqrt{\frac{d\sigma}{dt}} \quad t \simeq -\Delta^2$$


“amplitude”

challenges:

- coherent part drops fast with t ; statistical errors become large beyond $\sim 4^{\text{th}}$ minima
- accessible t range rather small ($|t| \lesssim 0.2 \text{ GeV}^2$) for coherent part

... so, does it work at all?

✓ yes, $|t| < 0.15 \text{ GeV}^2$ enough in eA to reconstruct Woods-Saxon potential used in simulation




exclusive processes and GPDs

another class of (related) processes for **parton imaging**

need to introduce concept of

generalized parton distributions (GPDs)



GPDs depend on:

- momentum transfer t
- resolution scale Q
- long. momentum before and after the scattering: x, ξ

= interference between different nucleon states (**not a probability**)



exclusive processes and GPDs

another class of (related) processes for **parton imaging**

need to introduce concept of



generalized parton distributions (GPDs)

GPDs depend on:


- momentum transfer t
- resolution scale Q
- long. momentum before and after the scattering: x, ξ

= interference between different nucleon states (**not a probability**)

appear in theoretical description of **exclusive processes**



deeply virtual Compton scattering (DVCS)



vector meson production


path to spatial imaging of partons through GPDs

recall: standard PDFs do not resolve transverse positions in the nucleon

fast moving nucleon turns into a 'pizza' but transverse size remains ≈ 1 fm

compelling questions


- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations



path to spatial imaging of partons through GPDs

recall: standard PDFs do not resolve transverse positions in the nucleon

fast moving nucleon turns into a 'pizza' but transverse size remains ≈ 1 fm



compelling questions

- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations

already discussed

$$f(x, k_T)$$

3-D

transv. mom. dep. PDF
semi-inclusive DIS

$$\int d^2 k_T$$

1-D


$$f(x)$$

parton densities

path to spatial imaging of partons through GPDs


recall: standard PDFs do not resolve transverse positions in the nucleon

fast moving nucleon turns into a 'pizza' but transverse size remains ≈ 1 fm



compelling questions


- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations



path to spatial imaging of partons through GPDs


recall: standard PDFs do not resolve transverse positions in the nucleon

fast moving nucleon turns into a 'pizza' but transverse size remains ≈ 1 fm



compelling questions


- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations



path to spatial imaging of partons through GPDs


recall: standard PDFs do not resolve transverse positions in the nucleon

fast moving nucleon turns into a 'pizza' but transverse size remains ≈ 1 fm



compelling questions


- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations



path to spatial imaging of partons through GPDs


recall: standard PDFs do not resolve transverse positions in the nucleon

fast moving nucleon turns into a 'pizza' but transverse size remains ≈ 1 fm



compelling questions


- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations



path to spatial imaging of partons through GPDs


recall: standard PDFs do not resolve transverse positions in the nucleon

fast moving nucleon turns into a 'pizza' but transverse size remains ≈ 1 fm



compelling questions


- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations



path to spatial imaging of partons through GPDs

recall: standard PDFs do not resolve transverse positions in the nucleon


fast moving nucleon turns into a 'pizza' but transverse size remains ≈ 1 fm



compelling questions

- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations


GPD's appear here



path to spatial imaging of partons through GPDs

recall: standard PDFs do not resolve transverse positions in the nucleon

fast moving nucleon turns into a 'pizza' but transverse size remains ≈ 1 fm




compelling questions

- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations

GPD's appear here


again, need Fourier transform
to get what we want



path to spatial imaging of partons through GPDs


recall: standard PDFs do not resolve transverse positions in the nucleon

fast moving nucleon turns into a 'pizza' but transverse size remains ≈ 1 fm



compelling questions


- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations



path to spatial imaging of partons through GPDs


recall: standard PDFs do not resolve transverse positions in the nucleon

fast moving nucleon turns into a 'pizza' but transverse size remains ≈ 1 fm



compelling questions

- how are quarks and gluons spatially distributed
- how do they move in the transverse plane
- do they orbit and do we have access to spin-orbit correlations



roadmap for transverse imaging of the proton

- ▶ obtain GPDs from global analysis of DVCS and vector meson data in ep scattering
slew of angular & polarization observables (+ use of positron beams) to disentangle H and E

4 GPDs per flavor, e.g.,

$$\int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p', s' | \bar{q}(-\frac{z}{2}) \mathcal{W} \gamma^+ q(\frac{z}{2}) | p, s \rangle_{z^+=0, \mathbf{z}=0}$$
$$= \quad \color{red}{H^q} \bar{u}(p', s') \gamma^+ u(p, s) + \color{red}{E^q} \bar{u}(p', s') \frac{i}{2m_p} \sigma^{+\alpha} (p' - p)_\alpha u(p, s)$$

roadmap for transverse imaging of the proton

- ▶ obtain GPDs from global analysis of DVCS and vector meson data in ep scattering
slew of angular & polarization observables (+ use of positron beams) to disentangle H and E

4 GPDs per flavor, e.g.,

$$\int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p', s' | \bar{q}(-\frac{z}{2}) \mathcal{W} \gamma^+ q(\frac{z}{2}) | p, s \rangle_{z^+=0, \mathbf{z}=0}$$
$$= H^q \bar{u}(p', s') \gamma^+ u(p, s) + E^q \bar{u}(p', s') \frac{i}{2m_p} \sigma^{+\alpha} (p' - p)_\alpha u(p, s)$$

recover PDFs in limit
 $s = s', \xi = 0, t = 0$

roadmap for transverse imaging of the proton

- ▶ obtain GPDs from global analysis of DVCS and vector meson data in ep scattering
slew of angular & polarization observables (+ use of positron beams) to disentangle H and E

4 GPDs per flavor, e.g.,

$$\int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p', s' | \bar{q}(-\frac{z}{2}) \mathcal{W} \gamma^+ q(\frac{z}{2}) | p, s \rangle_{z^+=0, \mathbf{z}=0}$$

$$= H^q \bar{u}(p', s') \gamma^+ u(p, s) + E^q \bar{u}(p', s') \frac{i}{2m_p} \sigma^{+\alpha} (p' - p)_\alpha u(p, s)$$

recover PDFs in limit
 $s = s', \xi = 0, t = 0$

no PDF limit; involves helicity flip
indicator of OAM; key part in Ji's sum rule

roadmap for transverse imaging of the proton

- ▶ obtain GPDs from global analysis of DVCS and vector meson data in ep scattering
slew of angular & polarization observables (+ use of positron beams) to disentangle H and E

4 GPDs per flavor, e.g.,

$$\int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p', s' | \bar{q}(-\frac{z}{2}) \mathcal{W} \gamma^+ q(\frac{z}{2}) | p, s \rangle_{z^+=0, \mathbf{z}=0}$$

$$= H^q \bar{u}(p', s') \gamma^+ u(p, s) + E^q \bar{u}(p', s') \frac{i}{2m_p} \sigma^{+\alpha} (p' - p)_\alpha u(p, s)$$


recover PDFs in limit
 $s = s', \xi = 0, t = 0$

no PDF limit; involves helicity flip
indicator of OAM; key part in Ji's sum rule


- ▶ perform Fourier transformation to obtain b-space image

e.g. $q(x, b^2) \simeq \int d^2\Delta e^{-ib\Delta} H^q(x, \xi = 0, t = -\Delta^2)$ where $\Delta = p' - p$

gives distribution of quarks with



- longitudinal momentum fraction x
- transverse distance b from proton center



roadmap for transverse imaging of the proton

- ▶ obtain GPDs from global analysis of DVCS and vector meson data in ep scattering
slew of angular & polarization observables (+ use of positron beams) to disentangle H and E

4 GPDs per flavor, e.g.,

$$\int \frac{dz^-}{4\pi} e^{ixP^+z^-} \langle p', s' | \bar{q}(-\frac{z}{2}) \mathcal{W} \gamma^+ q(\frac{z}{2}) | p, s \rangle_{z^+=0, \mathbf{z}=0}$$

$$= H^q \bar{u}(p', s') \gamma^+ u(p, s) + E^q \bar{u}(p', s') \frac{i}{2m_p} \sigma^{+\alpha} (p' - p)_\alpha u(p, s)$$


recover PDFs in limit
 $s = s', \xi = 0, t = 0$

no PDF limit; involves helicity flip
indicator of OAM; key part in Ji's sum rule


- ▶ perform Fourier transformation to obtain b-space image

e.g. $q(x, b^2) \simeq \int d^2\Delta e^{-ib\Delta} H^q(x, \xi = 0, t = -\Delta^2)$ where $\Delta = p' - p$

gives distribution of quarks with



- longitudinal momentum fraction x
- transverse distance b from proton center




challenges:

- need to resolve small distances in proton
- no diffractive pattern in accessible t range


→ need $d\sigma/dt$ in larger t -range than in eA

example: DVCS - what do we know?




- best understood and worked out theoretically
- x is integrated out in scattering amplitude
- ξ is related to usual Bjorken x of DIS: $\xi = x_{\text{Bj}} / (2 - x_{\text{Bj}})$
- large number of angular and polarization observables
- interferes with genuine QED “Bethe-Heitler” process

example: DVCS - what do we know?



- best understood and worked out theoretically
- x is integrated out in scattering amplitude
- ξ is related to usual Bjorken x of DIS: $\xi = x_{Bj} / (2 - x_{Bj})$
- large number of angular and polarization observables
- interferes with genuine QED “Bethe-Heitler” process



DVCS measurements
past-present-future



plenty of opportunities at an EIC
to map out gluon and sea-quark GPDs



DVCS - how well does it work out?

examples of simulated DVCS data including acceptance cuts and resolution effects





DVCS - how well does it work out?

examples of simulated DVCS data including acceptance cuts and resolution effects






Fourier




DVCS - how well does it work out?


examples of simulated DVCS data including acceptance cuts and resolution effects



Fourier






includes uncertainty from
different extrapolations for t




DVCS - how well does it work out?

examples of simulated DVCS data including acceptance cuts and resolution effects




includes uncertainty from
different extrapolations for t

Fourier




DVCS - what can we learn?



- ▶ unpolarized DVCS mainly sensitive to GPD H
- ▶ unknown GPD E from angular asymmetries with transversely polarized protons

different assumptions for GPD E

DVCS - what can we learn?




- ▶ unpolarized DVCS mainly sensitive to GPD H
- ▶ unknown GPD E from angular asymmetries with transversely polarized protons


different assumptions for GPD E

global QCD fits to data & Fourier transform to b space

quarks (unpolarized proton)




quarks (transversely pol. proton)




DVCS - what can we learn? - cont'd


quarks (unpolarized proton)



quarks (transversely pol. proton)




density of partons in transverse plane




DVCS - what can we learn? - cont'd


quarks (unpolarized proton)



quarks (transversely pol. proton)




density of partons in transverse plane




DVCS - what can we learn? - cont'd

quarks (unpolarized proton)



quarks (transversely pol. proton)



icing on the cake:


with GPDs H and E determined, one can access 'generalized form factors' by taking x moments, e.g.,

Ji's sum rule

$$J_{q,g} = \frac{1}{2} \int dx x [H^{q,g}(x, \xi, t \rightarrow 0) + E^{q,g}(x, \xi, t \rightarrow 0)]$$



total angular momentum of quarks and gluons

density of partons in transverse plane



gluon imaging through J/ψ production

- DVCS permits determination of gluon GPD through Q^2 evolution (similar to DIS)
- can be further improved by adding vector meson observables



take away message



an EIC offers many **unique opportunities** to greatly advance our understanding of the structure of nucleons and nuclei

precision studies of PDFs, TMDs, and GPDs will lead to the most comprehensive picture of the nucleon ever: its flavor, spin, and spatial structure

take away message



an EIC offers many **unique opportunities** to greatly advance our understanding of the structure of nucleons and nuclei

precision studies of PDFs, TMDs, and GPDs will lead to the most comprehensive picture of the nucleon ever: its flavor, spin, and spatial structure

requirements

- ▶ large enough c.m.s. energy to explore small x region
- ▶ sufficient luminosity for multi-dimensional binning, ...
- ▶ sufficient control of systematic uncertainties
- ▶ state-of-the-art detector systems, well integrated into IR



take away message



an EIC offers many **unique opportunities** to greatly advance our understanding of the structure of nucleons and nuclei

precision

entire science program uniquely tied to a future high-energy electron-ion collider

never been measured before & never without

TMDs, and GPDs

for comprehensive
the nucleon ever:
spin, and spatial structure

requirements

- ▶ large enough c.m.s. energy to explore small x region
- ▶ sufficient luminosity for multi-dimensional binning, ...
- ▶ sufficient control of systematic uncertainties
- ▶ state-of-the-art detector systems, well integrated into IR

